

Please read this leaflet in conjunction with the information provided in the dispensed Pregabalin packs.

Notes for your GP: Your patient has been prescribed pregabalin for neuropathic pain. Their dose has been titrated up to 150mg bd. If the medication has been helpful, please continue the prescription at 150mg bd. If the drug has proved ineffective, please tail off over two weeks or titrate up to 300mg bd if tolerated. Please prescribe one capsule bd at the final tolerated dose.

Extra notes:

Oxford Pain Consultants

Pregabalin (pre-GAB-a-lin)

50mg capsules

It is important that you keep this leaflet and show it to your pharmacist and GP

This information leaflet is designed to give you necessary information about this medicine. It will help you clearly understand why your doctor has prescribed it, benefits, possible side effects, how to take it and who to contact for more advice.

Oxford Pain Consultants
c/o Manor Hospital
Headington
Oxford
OX3 0PE
Tel: 01865 307734

What is Pregabalin (Lyrica)?

- Pregabalin is a new drug. It has been licensed for the treatment of nerve pain. It is different from over the counter painkillers, which might be why your Doctor has advised you take it.

How do I take it?

- The dose of Pregabalin will be different for different patients. The dose slowly increases. Please follow the table. Do **not** stop taking Pregabalin suddenly.

Please take Pregabalin as shown below.

| Day | AM | PM |
|-----|----------|----------|
| 1 | - | 1 x 50mg |
| 2 | - | 1 x 50mg |
| 3 | 1 x 50mg | 1 x 50mg |
| 4 | 1 x 50mg | 2 x 50mg |
| 5 | 2 x 50mg | 2 x 50mg |
| 6 | 2 x 50mg | 2 x 50mg |
| 7 | 2 x 50mg | 2 x 50mg |
| 8 | 2 x 50mg | 3 x 50mg |
| 9 | 2 x 50mg | 3 x 50mg |
| 10 | 3 x 50mg | 3 x 50mg |

What benefit can I expect?

- Nerve pain is difficult to treat. ‘Normal’ painkillers often don’t work. Pregabalin works by reducing the pain signals. You may find your pain level falls, which will help you to cope better.

What are the precautions?

- This medicine is usually well tolerated. It will add to the effects of alcohol or other drugs that can make you drowsy.
- Pregabalin may cause drowsiness, dizziness blurred vision, clumsiness, or difficulty in thinking.
- *Some patients may experience mood swings and changes to their behaviour. In the unlikely event symptoms such as increased anxiety and agitation should be reported to the GP.*
- Be careful how you react to this medicine before you drive a car or use heavy machinery
- Side effects are often related to the size of the dose and are usually temporary.
- You may need to reduce the dose to balance the benefits with any side effects
- Pregabalin is not addictive. However if you need to stop taking it, you should do so slowly. You can ask your Doctor about this. Some people feel sick if they stop it too quickly
- Pregabalin has not been tested in pregnancy.